

Measurement of the $E_{C.M.}=184$ keV resonance strength in the $^{26g}\text{Al}(p,\gamma)^{27}\text{Si}$ reaction

C. Ruiz,^{1,*} A. Parikh,^{2,†} J. José,^{3,4} L. Buchmann,¹ J. A. Caggiano,¹ A. A. Chen,⁵ J. A. Clark,² H. Crawford,⁶
B. Davids,¹ J. M. D'Auria,⁶ C. Davis,¹ C. Deibel,² L. Erikson,⁷ L. Fogarty,⁸ D. Frekers,⁹ U. Greife,⁷
D. A. Hutcheon,¹ M. Huyse,¹⁰ C. Jewett,⁷ A. M. Laird,¹¹ R. Lewis,² P. Mumby-Croft,¹¹ A. Olin,¹ D. F. Ottewell,¹
C. V. Ouellet,⁵ P. Parker,² J. Pearson,⁵ G. Ruprecht,¹ M. Trinczek,¹ C. Vockenhuber,¹ and C. Wrede²

¹*TRIUMF, Vancouver, BC V6T 2A3, Canada*

²*Wright Nuclear Structure Laboratory, Yale University, New Haven, CT 06520-8124, USA*

³*Dept. de Física i Enginyeria Nuclear, Universitat Politècnica de Catalunya, Barcelona, Spain*

⁴*Institut d'Estudis Espacials de Catalunya (IEEC), Barcelona, Spain*

⁵*McMaster University, Hamilton, ON L8S 4L8, Canada*

⁶*Simon Fraser University, Burnaby, BC V5A 1S6, Canada*

⁷*Dept. of Physics, Colorado School of Mines, Golden, CO 80401, USA*

⁸*National University of Ireland, Maynooth, Co. Kildare, Ireland*

⁹*Institut für Kernphysik, Westfälische Wilhelms-Universität Münster, 48149 Münster, Germany*

¹⁰*Katholieke Universiteit Leuven, 3000 Leuven, Belgium*

¹¹*Dept. of Physics, University of York, York YO10 5DD, UK*

(Dated: April 13, 2006)

The strength of the $E_{C.M.}=184$ keV resonance in the $^{26g}\text{Al}(p,\gamma)^{27}\text{Si}$ reaction has been measured in inverse kinematics using the DRAGON recoil separator at TRIUMF's ISAC facility. We measure a value of $\omega\gamma = 35 \pm 7$ μeV and a resonance energy of $E_{C.M.}=184\pm 1$ keV, consistent with p-wave proton capture into the 7652(3) keV state in ^{27}Si , and discuss the implications of these values for ^{26g}Al nucleosynthesis in typical oxygen-neon white-dwarf novae.

PACS numbers: 24.30.-v, 25.40.Lw, 26.50.+x

Keywords: ^{26}Al , novae, radioactive ion beams, γ -ray astronomy

Introduction - The recent detection of decaying ^{26g}Al ($t_{1/2}=(7.1\pm 0.2)\times 10^5$ yr) via its characteristic 1.809 MeV γ -ray by the RHESSI and INTEGRAL satellites has furthered our understanding of the production sites of this radioisotope [1] [2]. The COMPTEL all-sky map of the 1.809 MeV line [3] points to young, high-mass progenitors such as core collapse supernovae (CCSN) and Wolf-Rayet stars [4]. Though previous studies suggested that the measured $2.8\pm 0.8M_{\odot}$ [5] of ^{26g}Al in the Galaxy could have been entirely produced in CCSN [6], these new results have suggested that CCSN may be a much less dominant component, and that other sources must contribute, thought chiefly to be Wolf-Rayet stars [7].

However, classical novae are one potential source of ^{26g}Al and it has been shown that up to $0.4M_{\odot}$ of the Galactic abundance could have been produced in these sites [8]. Of particular importance to the calculation of nova-synthesized ^{26g}Al abundances are the $^{25}\text{Al}(p,\gamma)^{26}\text{Si}$ and $^{26g}\text{Al}(p,\gamma)^{27}\text{Si}$ reaction rates, the former being the most uncertain. The $^{26g}\text{Al}(p,\gamma)^{27}\text{Si}$ reaction rate at typical oxygen-neon white-dwarf nova (ONeWD) temperatures is dominated by a single resonance around $E_{C.M.}=188$ keV [9, 10].

An unpublished measurement of this resonance

strength in normal kinematics yielded a value of 55 ± 9 μeV [9]. The final abundance of ^{26g}Al synthesised in a typical ONeWD hydrodynamic calculation is somewhat sensitive to this strength. Due to the long lifetime of ^{26g}Al , space-based γ -ray observatories such as INTEGRAL are unable to detect it from individual sources. Therefore the likely primary progenitors can only be inferred from the Galactic ^{26g}Al distribution. However, with a firm understanding of the $^{26g}\text{Al}(p,\gamma)^{27}\text{Si}$ rate we can infer solid upper limits for the nova contribution to Galactic ^{26g}Al as a secondary source.

Experimental Method - This measurement was performed using the DRAGON recoil separator in the ISAC radioactive ion beam facility at TRIUMF. A high-power SiC target [11] was bombarded with up to 70 μA of 500 MeV protons from TRIUMF's sector focussing cyclotron, producing radioactive ^{26g}Al which then diffused out of the target and into a rhenium surface-ionization tube. An enhancement to the surface ionization was provided using on-line resonant-laser ionization within the tube [12]. $A=26$ products were separated using a high resolution mass separator. This beam was injected into a radio-frequency quadrupole accelerator (RFQ) for initial acceleration up to 0.15 A MeV and stripped to a higher charge state using a thin carbon foil before being injected into a continuously-variable energy drift-tube LINAC (DTL) which allowed acceleration between 0.15-1.8 A MeV [13]. The beam was delivered in bunches separated by 86 ns in time with a bunch width of less than two nanoseconds

*Electronic address: ruiz@triumf.ca; Also at: Simon Fraser University, Burnaby, BC V5A 1S6, Canada

†Electronic address: anuj.parikh@yale.edu

FWHM and an energy spread of typically 1% FWHM at 0.2 A MeV.

The DRAGON facility [14, 15] consists of a windowless hydrogen-recirculating gas-target surrounded by an array of 30 bismuth germanate (BGO) detectors, and a two-stage electromagnetic recoil separator. Each stage includes a dipole magnet and an electrostatic dipole unit. The total separator length from the centre of the gas cell to the final m/q focus is 20.42 m, with a double-sided silicon strip detector (DSSD) placed 65 cm downstream from the focus. A collimated silicon surface barrier detector placed at 30° from the centre of the gas cell was used to detect protons elastically scattered by the incoming beam during the run as a means of normalisation.

The separator was set to transmit charge-state 4^+ silicon recoils from the reaction $^{26g}\text{Al}(p,\gamma)^{27}\text{Si}$, which were then detected using the DSSD in coincidence with prompt γ -rays detected at the BGO array. The majority of unreacted 4^+ beam ended up impinging on a slit at the m/q focus after the first electric dipole, but a small fraction (1×10^{-9}) were transmitted to the DSSD. These “leaky-beam” particles, in random coincidence with background radiation in the BGO array, constituted a significant background under the true coincidence peak in a time of flight spectrum (figure 1). These ions had slightly higher energy than the ^{27}Si recoils in the DSSD energy spectrum. The BGO background came from ^{26}Na in the beam and from room radioactivity.

The contaminants in the beam were primarily radioactive ^{26}Na ($t_{1/2}=1.07$ s) and ^{26m}Al ($t_{1/2}=6.345$ s) created in the SiC target. The level of these contaminants in the beam was monitored during the run; the ^{26}Na via its characteristic 1809 keV γ -ray from beta-decay to the first excited state of ^{26}Mg using a high-purity Germanium detector; the ^{26m}Al via paired 511 keV γ -rays created from electron-positron annihilation caused by β^+ decays of this isomer at the mass slits. The ^{26}Na contaminant was the major source of background in the experiment due to the small amount of beam implanted in the entrance aperture to the gas cell causing a high rate in the BGO array leading to random coincidences. With a combination of a mechanical iris upstream of the gas cell (designed to remove the source of beam decay away from the detectors) and some fine tuning of the mass separator optics, we were able to reduce the level of ^{26}Na in the beam from 1:32,000 to 1:337,000, with a contribution to the BGO rate equal to that of room radioactivity. The level of ^{26m}Al contaminant remained at around 1:30,000 during the run.

During the run, a gas target pressure of 6 Torr was maintained to within $\pm 1.6\%$. Initially, a beam energy of 5.226 MeV ($E_{C.M.}=0.195$ MeV) was chosen to place the resonance at the centre of the gas target and over 179 hours of data were taken at an average intensity of 2.5×10^9 s $^{-1}$ at this energy, although peak intensities of over 5×10^9 s $^{-1}$ were achieved. Some 49 hours of data were taken at an energy of 5.122 MeV ($E_{C.M.}=0.191$ MeV), and an “off-resonance” background run was taken

at 5.850 MeV ($E_{C.M.}=0.218$ MeV) for 30 hours. In addition, a beam of ^{28}Si was produced in the ISAC off-line microwave ion-source by extracting $A=31$ ($^{28}\text{SiH}_3^+$) molecules into the RFQ and selecting $^{28}\text{Si}^{5+}$ for acceleration through the DTL. This beam was used to study the charge state distribution of silicon ions exiting the gas target at different pressures in order to account for charge states not measured in the experiment.

Analysis - From the coincident gamma-ray/heavy-ion data for each set of runs at a particular energy, true recoil events were identified by their time-of-flight through the separator. Since most random coincidences were caused by γ -rays from decaying ^{26}Na atoms upstream of the gas target and leaky beam particles at the DSSD, these were uncorrelated in time and resulted in a flat background in the separator time-of-flight spectrum. The recoils protrude from the flat time-of-flight background as a tightly-bunched peak, and are also peaked at lower energy than the leaky beam in the DSSD spectrum, as can be seen in figure 1.

Candidate $^{26}\text{Al}(p,\gamma)^{27}\text{Si}$ events were selected by a narrow window (200 ns) on time-of-flight (TOF) set around the coincidence peak. The number of background events in the narrow window was estimated from the number of events in a much wider window in a randoms-only region of the TOF spectrum. The background-subtracted numbers for the 5.226 MeV and 5.122 MeV runs were: 119 ± 14 recoils and 28 ± 6 recoils respectively. The number for the off-resonance run at 5.850 MeV was derived using the Feldman-Cousins prescription for non-Poissonian low-statistics data [16] leading to a limit of < 3.72 recoils at 90% confidence level.

After the subtraction of measured contaminants was made, the normalisation procedure described above resulted in values for the number of incident ions of 1.5×10^{15} (5.226 MeV runs), 3.5×10^{14} (5.122 MeV runs) and 2.9×10^{14} (5.850 MeV runs). The absolute reaction yields ([reactions/incident ion]) for these same runs were: $(2.5 \pm 0.5) \times 10^{-13}$, $(2.6 \pm 0.7) \times 10^{-13}$ and $< 4.0 \times 10^{-14}$ respectively.

The detected yields were corrected for the measured charge-state fraction for 4^+ recoils exiting the gas target ($\eta_{Si4+} = 0.42 \pm 0.02$), the separator acceptance ($\eta_{sep} = 0.98 \pm 0.02$), the efficiency of the BGO array ($\eta_{bgo} = 0.76 \pm 0.1$) and the previously known DSSD efficiency ($\eta_{DSSD} = 0.97 \pm 0.01$). η_{sep} was found by simulating the reaction in a full Monte Carlo implementation of the DRAGON recoil separator in GEANT3, including interaction of the γ -rays with the BGO array and the ion-optical tracking of charged particles through the separator. Uncertainties in the beam energy, energy spread, position and direction within the gas cell were included in the simulation. The maximum cone-angle of recoils from the reaction is around 15 mrad, and consequently a very small fraction of recoils are close to the acceptance design limit of the separator when factoring in beam emittance. As a result, a small fraction of recoils do not

make it through the system. η_{bgo} was determined using the GEANT3 simulation also [17], taking into account the effect of hardware thresholds and unknown γ -ray angular distributions. The sensitivity of the efficiency to the γ -ray branching ratios from the 7652 keV state was also investigated, and included in the experimental uncertainty.

Stopping powers for the different beam energies in the H_2 gas were measured by taking the difference between the field strengths of the first-stage magnetic dipole required to center the beam at an energy-dispersed focus with and without gas in the target. This procedure was calibrated using some well-known narrow radiative capture resonances [15].

Resonance strengths were calculated using the formula

$$\omega\gamma = \frac{2\epsilon Y}{\lambda^2} \frac{M_H}{M_{Al} + M_H} \quad (1)$$

where λ is the de-Broglie wavelength in the C.M. system, ϵ is the beam energy loss per target atom per unit area, Y is the reaction yield, and M_H, M_{Al} are the masses of the target and projectile respectively.

The reaction yield is given by


$$Y = N_{det}/(N_{inc}\eta_{bgo}\eta_{sep}\eta_{Si4+}\eta_{DSSD}) \quad (2)$$

where N_{det} and N_{inc} are the total number of detected recoils and incident ions respectively. Table I shows the resulting resonance strengths and their associated errors.


The location of the resonant capture within the extended gas target was deduced from the pattern of hits in the BGO array, shown in figure 2. The associated error on this measurement is ± 9 mm. The energy of the resonance was calculated by correcting the incident beam energy for energy loss before reaching the resonance position. The calculated resonance energy is 184 ± 1 keV.

The resonance strength obtained in this work is only 64% of the unpublished value of ref. [9], while the resonance energy is measured to be 2% smaller. This has the effect of reducing the reaction rate over the region of the Gamow window for this resonance by nearly a factor of 1.2. A slower reaction rate means that for a given temperature, more ^{26}Al will survive a nova explosion.

Astrophysical implications: synthesis of ^{26}Al in nova outbursts - An analysis of the impact of the new $^{26g}Al(p,\gamma)^{27}Si$ rate on the synthesis of ^{26g}Al in novae has been performed. We will not attempt to reach final conclusions on the precise contribution of novae to the Galactic ^{26g}Al levels because of the uncertainties facing such calculations (including the important role played by nuclear uncertainties affecting the $^{25}Al(p,\gamma)^{26}Si$ rate, a channel that bypasses the synthesis of ^{26g}Al in novae through $^{25}Al(p,\gamma)^{26}Si(\beta^+)^{26m}Al(\beta^+)^{26}Mg$; see [18] for details). Current estimates suggest that novae contribute less than 20% of the Galactic ^{26g}Al abundances [8]. However, we can outline the extent to which this new rate affects ^{26g}Al synthesis in novae by computing a



(a)



(b)

FIG. 1: (a) Separator time-of-flight for coincident gamma-ray/heavy-ion events versus detected particle energy for the 5.122 MeV run. The true ^{27}Si recoils are bunched tightly in time, and are peaked at lower energy than the randomly-coincident 'leaky' beam particles. (b) Projection of (a) onto the time-of-flight axis showing the true recoil peak.

representative case. Hence we have computed new simulations of nova outbursts, assuming an accreting ONe white dwarf of $1.25 M_{\odot}$, from the onset of accretion up to the explosion and ejection stages, by means of a spherically symmetric, implicit, hydrodynamic code, in Lagrangian formulation [19]. We have compared the mean ^{26}Al yields in the ejecta when two different prescriptions for the $^{26g}Al(p,\gamma)$ rate are adopted: one corresponding to the unpublished (Vogelaar [9]) rate and a second with the rate presented in this paper.

As a result of the lower strength associated with the $E_{C.M.} = 188$ keV resonance, the net reduction of the overall $^{26g}Al(p,\gamma)^{27}Si$ rate favors the synthesis of ^{26g}Al

Percentage contribution to error

E_{beam} (MeV)	$\Delta\epsilon$	$\Delta\eta_{DSSD}$	$\Delta\eta_{BGO}$	$\Delta\eta_{sep}$	$\Delta\eta_{Si4+}$	ΔN	Tot. sys. error	$\omega\gamma$ (μeV)
5.226	5%	1%	13%	2%	5%	3%	15%	$35 \pm 5_{sys} \pm 4_{stat}$
5.122	5%	1%	13%	2%	5%	8%	17%	$36 \pm 6_{sys} \pm 8_{stat}$

TABLE I: Table showing the percentage contributions to the systematic error for the parameters which go into the calculation of the resonance strength, $\omega\gamma$, for the 5.226 MeV and 5.122 MeV runs. Also tabulated are the obtained resonance strengths and associated errors (see text for definitions).




FIG. 2: Plot of z-coordinate of the BGO detector struck by the highest energy γ -ray coincident with a heavy-ion detection for the 5.122 MeV runs, showing the distribution of hits peaked at the center of the array.

in nova outbursts. The results obtained from the two hydrodynamic simulations lead to mean ^{26g}Al yields in

the ejecta of 6.1×10^{-4} by mass, when Vogelaar's values for the strength and energy of the 184 keV resonance are adopted (cf., $55\mu\text{eV}$ and 188 keV), or 7.4×10^{-4} , when the present values are used. This represents an increase of $\sim 20\%$ of the overall ^{26g}Al yield, for this particular model, representative of a neon-type nova. The importance of the results obtained from these hydrodynamics simulations is twofold: first, it confirms that classical novae are likely sites for the synthesis of a fraction of the Galactic ^{26g}Al ; and second, the moderate increase in the final ^{26g}Al yield is compatible with the current paradigm for the origin of the Galactic ^{26g}Al , namely young massive star progenitors, with the likely contribution of secondary sources such as novae and AGB stars.

The authors would like to acknowledge funding from the National Research Council of Canada (NRC), the National Science and Engineering Research Council (NSERC) and US Dept. of Energy grants DE-FG02-91ER-40609, DE-FG02-93ER-40789. We also gratefully acknowledge the invaluable assistance in beam production from P. Bricault, M. Domsbys, J. Lassen and team, K. Wendt (U. Mainz), R. E. Laxdal and M. Marchetto.

-
- [1] D. M. Smith, *Astrophysical Journal* **589**, L55 (2003).
[2] J. Knödseder, *New Astronomy Reviews* **48**, 189 (2004).
[3] R. Diehl et al., *Astronomy & Astrophysics* **298**, 445 (1995).
[4] N. Prantzos and R. Diehl, *Physics Reports* **267**, 1 (1996).
[5] R. Diehl et al., *Nature* **439**, 54 (2006).
[6] F. X. Timmes et al., *Astrophysical Journal* **449**, 204 (1995).
[7] N. Prantzos, *Astronomy & Astrophysics* **420**, 1033 (2004).
[8] J. José, M. Hernanz, and A. Coc, *Astrophysical Journal* **479**, L55 (1997).
[9] R. B. Vogelaar, Ph.D. thesis, California Institute of Technology (1989).
[10] R. B. Vogelaar et al., *Physical Review C* **53**, 1945 (1996).
[11] P. Bricault et al., *Nuc. Inst. Meth. B* **204**, 319 (2003).
[12] J. Lassen et al., *Hyperfine Interactions* DOI: **10.1007/s10751-005-9212-2** (2006).
[13] R. E. Laxdal, *Nuc. Inst. Meth. B* **204**, 400 (2004).
[14] D. A. Hutcheon et al., *Nuc. Inst. Meth. A* **498**, 190 (2003).
[15] S. Engel et al., *Nuc. Inst. Meth. A* **553**, 491 (2005).
[16] G. J. Feldman and R. D. Cousins, *Physical Review D* **57**, 3873 (1998).
[17] D. G. Gigliotti, J. G. Rogers, and A. H. Hussein, *Nuc. Inst. Meth. B* **204**, 671 (2003).
[18] J. José, A. Coc, and M. Hernanz, *Astrophysical Journal* **520**, 347 (1999).
[19] J. José and M. Hernanz, *Astrophysical Journal* **494**, 680 (1998).